

Child Immunization Schedule

Vaccine	Recommended At
Hepatitis B/HepB ¹	Birth-2 months
	1-4 months
	6-18 months
Rotavirus/RV ²	2 months
	4 months
	6 months
Diphtheria, Tetanus, Pertussis/DTaP	2 months
	4 months
	6 months
	15-18 months
	4-6 years
Inactivated Polio/IPV	2 months
	4 months
	6-18 months
	4-6 years
<i>H. influenzae</i> type b/Hib	2 months
	4 months
	6 months
	12-15 months

Vaccine	Recommended At
Pneumococcal/PCV (pneumonia, meningitis)	2 months
	4 months
	6 months
	12-15 months
Influenza ³	6 months-18 years
Measles, Mumps, Rubella/MMR	12-15 months
	4-6 years
Varicella (chickenpox)	12-15 months
	4-6 years
Hepatitis A/HepA	12-23 months
Tetanus, Diphtheria, Pertussis/Tdap ⁴	11-12 years
Meningococcal/MCV (meningitis)	11-12 years
Human Papillomavirus/HPV ⁵	11-12 years

If your child misses a recommended dose of a vaccine, ask your doctor about catch-up doses.

¹ HepB: All infants should receive the first dose within 12 hours of birth. The first dose may also be given by age 2 months if the mother is HBsAg-negative.

² Rotavirus: If the vaccine given at ages 2 and 4 months is Rotarix[®], a dose at 6 months is not needed.

³ Influenza: Seasonal flu—yearly between 6 months-18 years. H1N1 flu—check with your local health department.

⁴ Tdap: For children who have received the recommended childhood DTaP series and have not received a Td booster dose.

⁵ HPV: Available for females and males, given in 3 doses.